

NO-A178 939

VIBRATIONAL MOTIONS OF BUCKMINSTERFULLERENE(U) STATE
UNIV OF NEW YORK AT BUFFALO DEPT OF CHEMISTRY
Z C MU ET AL. MAR 87 UBUFFALO/DC/87/TR-34

1/1


UNCLASSIFIED

NO0014-86-K-0043

F/G 7/4

NL





AD-A178 939

OFFICE OF NAVAL RESEARCH

Contract N00014-K-0043

R & T Code 413f0001---01

TECHNICAL REPORT No. 34

Vibrational Motions of Buckminsterfullerene

by

Z. C. Wu, Daniel Jelski and Thomas F. George

Prepared for Publication

in

Chemical Physics Letters

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

March 1987

Reproduction in whole or in part is permitted for any purpose of
the United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
APR 07 1987
S D

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/87/TR-34			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
11. TITLE Vibrational Motions of Buckminsterfullerene					
12. PERSONAL AUTHOR(S) Z. C. Wu, Daniel Jelski and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) March 1987	
				15. PAGE COUNT 14	
16. SUPPLEMENTARY NOTATION Prepared for Publication in Chemical Physics Letters					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	BUCKMINSTERFULLERENE, NON-CARTESIAN COORDINATES		
			CARBON (60-ATOM) CLUSTER; FOUR FORCE CONSTANTS		
			COMPLETE VIBRATIONAL SPECTRUM, 180 BY 180 MATRIX		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A non-Cartesian coordinate system is developed which permits the vibrational motions of Buckminsterfullerene (Bucky ball) to be expressed in terms of four force constants. A 180 x 180 matrix is then derived which, when diagonalized, yields the complete vibrational spectrum. These results are compared with those obtained previously via a MNDO calculation.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

VIBRATIONAL MOTIONS OF BUCKMINSTERFULLERENE

Z. C. Wu, Daniel A. Jelski and Thomas F. George

Departments of Physics & Astronomy and Chemistry
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

Abstract

A non-Cartesian coordinate system is developed which permits the vibrational motions of Buckminsterfullerene (Bucky ball) to be expressed in terms of four force constants. A 180×180 matrix is then derived which, when diagonalized, yields the complete vibrational spectrum. These results are compared with those obtained previously via a MNDO calculation.



Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

I. Introduction

The purpose of this paper is to investigate the vibrational motions of Buckminsterfullerene (Bucky ball) in a way that lends itself toward future application. Hence simplicity, both theoretical and practical, is an important consideration. To this end we can take advantage of the symmetry of the species, and we shall derive a relatively simple procedure for calculating the vibrational normal frequencies. Furthermore, our method lends itself to physical interpretation, and we shall be able to make some statements about the nature of Bucky ball. The experimental literature on Bucky Ball is well known.¹⁻⁵ Theoretically much work has been dedicated to determining the most stable structure of C_{60} clusters,⁶⁻¹⁰ and to determining some of their optical properties.¹¹ We continue this work here by assuming that Bucky ball exists and is relatively stable.

In essence our calculation can be described as follows. We express the position of each atom in local coordinates, choosing our coordinate system (non-Cartesian) in a way that simplifies the calculation. This permits us to express the equation of motion of each atom in terms of four force constants, and since all atoms are identical we immediately derive an equation of motion for the entire system. The fundamental principles behind this calculation are found in Ref. 12. While we must input the force constants into our calculation (this is a disadvantage), we generate the correct number of modes, the correct point group properties, and, at least qualitatively, the correct frequencies. This requires only the diagonalization of one 180×180 matrix, a procedure which requires only about three minutes CPU on a VAX. Furthermore, since we can vary the force

constants and since we can compare our results with those of Stanton,¹³ we are able to make some predictions about the structure of Bucky ball as compared to benzene. In the next section we derive our method for calculating the normal mode frequencies, and in Section III we compare our results with those of Stanton¹³ and discuss the implications.

II. Theory

The structure of Buckminsterfullerene, illustrated in Fig. 1, is that of a truncated icosahedron with sixty vertices, twenty hexagonal faces and twelve pentagonal faces. A carbon atom occupies each vertex. The bonds separating a hexagon from a pentagon are found to be more "single" than bonds separating two hexagons.^{6,8} The bond lengths are given as 1.41 Å and 1.54 Å, respectively.⁸ Since each vertex is at the intersection of two hexagonal and one pentagonal face, it follows that it is joined by one "double" bond and two "single" bonds. This implies that the motion of an atom around its equilibrium position can be decomposed along the three non-coplanar directions of the adjacent bonds (Fig. 2). We can use x_i , y_i and z_i to denote the deviation of atom i from its equilibrium position in these directions (we caution the reader not to confuse our notation with Cartesian coordinates, since ours are not Cartesian). The angle between two adjacent single bonds is $\frac{3\pi}{5}$ and that between a double bond and an adjacent single bond is $\frac{2\pi}{3}$. One can work out the angle between a double bond and its adjacent pentagonal face as $\Psi = \cos^{-1}\left(\frac{1}{2\cos\frac{3\pi}{5}}\right)$. The angle between two adjacent hexagonal faces is $\Phi = \cos^{-1}\left[\frac{8}{3}\left(\sin\frac{3\pi}{10}\right)^2 - 1\right]$, and the angle between adjacent hexagonal and pentagonal faces is

$$\theta = \frac{\pi - \phi}{2} - \psi.$$

The variations of the bond length a_{ij} , b_{ik} and c_{il} (see Fig. 2) due to the motion of the atoms are, to first order,

$$\begin{aligned}\delta a_{ij} &= (x_i - \frac{1}{2}y_i - \frac{1}{2}z_i) + (x_j - \frac{1}{2}y_j - \frac{1}{2}z_j), \\ \delta b_{ik} &= (y_i - \frac{1}{2}x_i - \cos\frac{2\pi}{5}z_i) + (z_k - \frac{1}{2}x_k - \cos\frac{2\pi}{5}y_k), \\ \delta c_{il} &= (z_i - \frac{1}{2}x_i - \cos\frac{2\pi}{5}y_i) + (y_l - \frac{1}{2}x_l - \cos\frac{2\pi}{5}z_l).\end{aligned}\quad (1)$$

Also, the variations of the angles between bonds α , β and γ are

$$\begin{aligned}\delta\alpha_i &= \frac{1}{L}(-\sin\frac{\pi}{3}\cos\phi y_j + \sin\frac{\pi}{3}z_j + \sin\frac{\pi}{3}x_k - \sin\frac{2\pi}{5}\cos\theta y_k \\ &\quad - \sin\frac{\pi}{3}y_i + \sin\frac{\pi}{3}\cos\phi z_i - \sin\frac{\pi}{3}x_i + \sin\frac{2\pi}{5}\cos\theta z_i), \\ \delta\beta_i &= \frac{1}{L}(-\sin\frac{\pi}{3}\cos\phi z_j + \sin\frac{\pi}{3}y_j + \sin\frac{\pi}{3}x_l - \sin\frac{2\pi}{5}\cos\theta z_l \\ &\quad - \sin\frac{\pi}{3}z_i + \sin\frac{\pi}{3}\cos\phi y_i - \sin\frac{\pi}{3}x_i + \sin\frac{2\pi}{5}\cos\theta y_i), \\ \delta\gamma_i &= \frac{1}{L}(\sin\frac{2\pi}{5}y_k - \sin\frac{\pi}{3}\cos\theta x_k + \sin\frac{2\pi}{5}z_l - \sin\frac{\pi}{3}\cos\theta x_l \\ &\quad + \sin\frac{\pi}{3}\cos\theta x_i - \sin\frac{2\pi}{5}y_i + \sin\frac{\pi}{3}\cos\theta x_i - \sin\frac{2\pi}{5}z_i),\end{aligned}\quad (2)$$

where L refers to the bond lengths which are assumed equal.

The kinetic energy of each atom is

$$T_1 = \frac{m}{2} \left(\dot{x}_1 - \frac{1}{2}\dot{y}_1 - \frac{1}{2}\dot{z}_1 \right)^2 + \left(\dot{y}_1 \cos \frac{\pi}{5} - \dot{z}_1 \cos \frac{\pi}{5} \right)^2 + \left(\dot{y}_1 \cos \frac{3\pi}{10} \sin \varphi + \dot{z}_1 \cos \frac{3\pi}{10} \sin \varphi \right)^2, \quad (3)$$

where m is the mass of each atom. It is assumed that the atoms undergo harmonic oscillations around their equilibrium positions, where the potential is due to the variations of the bond lengths and the angles between them. If atoms j , k and l represent nearest neighbors of atom i , as shown in Fig. 2, then the Lagrangian for this system takes the form

$$L = T - V = \sum_i T_i - \sum_{i>j} \frac{1}{2} k_1 \delta a_{ij}^2 - \sum_{i>k} \frac{1}{2} k_2 \delta b_{ik}^2 - \sum_{i>l} \frac{1}{2} k_2 \delta c_{il}^2 - \sum_i \frac{1}{2} [k_3 (\delta \alpha_i^2 + \delta \beta_i^2) + k_4 \delta \gamma_i^2], \quad (4)$$

where k_1 , k_2 , k_3 and k_4 are all force constants. Equation (4) can be rewritten in the compact form

$$L = \frac{1}{2} \sum_{m,n=1}^{180} (T_{mn} X^m X^n - V_{mn} X^m X^n), \quad (5)$$

and the appropriate Euler-Lagrange equation is

$$\ddot{X}^n = - \sum_{m,l=1}^{180} (T^{-1})^{mn} V_{ml} X^l. \quad (6)$$

By diagonalizing the 180×180 matrix $T^{-1}V$, we get the squares of the eigenfrequencies along with their associated eigenvectors.

Several cases are worth discussing:

- (a) $k_2 = k_3 = k_4 = 0$. Each double bond with force constant k_1 becomes independent. Then one obtains only one non-zero eigenfrequency, $(k_1/m)^{1/2}$, with a 30-fold degeneracy.
- (b) $k_1 = k_3 = k_4 = 0$. The system reduces to 12 independent pentagons. For each pentagon the eigenfrequencies are (with degeneracies) 0.89401 (1), 1.00029 (2) and 1.15166 (2), with $k_2/m = 1$. All other eigenfrequencies are zero.
- (c) $k_3 = k_4 = 0$. One obtains 90 non-zero eigenfrequencies corresponding to the 90 bonds. Since there is no force on the angles, Bucky ball can be squished in $90 - 6 = 84$ different ways. The remaining 6 degrees of freedom are due to translation and rotation.
- (d) In the general case, there are always 6 zero eigenfrequencies corresponding to global translational and rotational motions. There are also three non-degenerate modes.

III. Results and Discussion

Table I contains 180 eigenvalues representing the vibrational frequencies of Buckminsterfullerene. The values of $k_1 - k_4$ are those that best seem to fit the data of Stanton,¹³ which was calculated by the MNDO method with the gradients evaluated at six points around an atom. The best fit was determined by matching the highest and lowest frequencies⁶ and also by matching the three non-degenerate frequencies. In addition, we have insisted that the ratio between k_1 and k_2 be approximately equal to the

inverse ratio of the bond lengths. Since at this point we have no independent way of determining k_4 , we have insisted that it equal k_3 .

For benzene the appropriate values would be $k_1 = k_2 = 7.62 \times 10^5$ dynes/cm and $k_3 = .667 \times 10^5$ dynes/cm.¹⁴ Comparing these values with those given in Table I, we see that the force constants for Bucky ball are considerably higher than for benzene. This is physically reasonable since the stretching of a bond in Bucky ball involves a dislocation of the entire molecule, and hence greater resistance. Similarly we expect the stretching constant to be larger.

It is also instructive to look at the three non-degenerate modes. Stanton¹³ has calculated their frequencies (in cm^{-1}) as 611 (for the A_g mode), 973 (A_u) and 1667 (A_g). Our calculation reveals frequencies at 548, 970, and 1627 cm^{-1} , respectively. We have found that the two A_g modes depend only on constants k_1 and k_2 . The first is a "breathing" motion whereas the second involves oscillation between the "single" and "double" bonds. In both cases angles can be expected to play no role. The A_u motion, on the other hand, involves a twisting (angle-dependent) motion of the pentagons. When k_3 and k_4 are zero, the eigenfrequency of this motion becomes zero. All other modes must be degenerate according to the point group of Buckminsterfullerene.¹³

Our method is well suited to elucidating this sort of physical description of the motions since the k 's have obvious physical significance. This is a big advantage of this calculation, along with the numerical simplicity. We intend to use this method to further investigate the vibrational spectrum of Buckminsterfullerene.

Acknowledgments

We gratefully acknowledge the assistance of Dr. Richard E. Stanton for sharing with us some unpublished research results. We also thank him for useful discussions. This research was supported by the Office of Naval Research and the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints notwithstanding any copyright notation hereon.

References


1. H.W. Kroto, J.R. Heath, S.C. O'Brien, R.F. Curl and R.E. Smalley, *Nature* 318, 162 (1985). For a review of the experimental techniques involved, see A. Kaldor, D.M. Cox, D.J. Trevor and M.R. Zakin, *Z. Phys. D* 3, 195 (1986).
2. Q.L. Zhang, S.C. O'Brien, J.R. Heath, Y. Liu, H.W. Kroto and R.E. Smalley, *J. Phys. Chem.* 90, 525 (1986).
3. A. O'Keefe, M.M. Ross and A.P. Baronavski, *Chem. Phys. Lett.* 130, 17 (1986).
4. M.Y. Hahn, E.C. Honea, A. J. Paguia, K.E. Schriver, A.M. Camarena and R.L. Whetten, *Chem. Phys. Lett.* 130, 12 (1986).
5. S.C. O'Brien, J.R. Heath, H.W. Kroto, R.F. Curl and R.E. Smalley, *Chem. Phys. Lett.* 132, 99 (1986).
6. M.D. Newton and R.E. Stanton, *J. Am. Chem. Soc.* 108, 2469 (1986).
7. T.G. Schmalz, W.A. Seitz, D.J. Klein and G.E. Hite, *Chem. Phys. Lett.* 130, 203 (1986).
8. S. Satpathy, *Chem. Phys. Lett.* 130, 545 (1986).
9. R.C. Haddon, L.E. Brus and K. Raghavachari, *Chem. Phys. Lett.* 131, 165 (1986).
10. P.W. Fowler, *Chem. Phys. Lett.* 131, 444 (1986).
11. D.A. Jelski and T.F. George, *J. Phys. Chem.*, submitted.
12. E.B. Wilson, J.C. Decius and P.C. Cross, Molecular Vibrations (McGraw-Hill, New York, 1955).
13. R.E. Stanton, private communication.
14. B.L. Crawford and F.A. Miller, *J. Chem. Phys.* 17, 249 (1949).


Frequency	Symmetry	Frequency	Symmetry
0.0	-	272.0	Hg
354.7	Hu	361.8	F2u
373.7	Gu	427.9	Hg
455.8	Gg	491.3	Flu
491.6	Hu	525.2	F2g
530.2	Gg	547.6	Ag
550.9	Flu	552.4	Hg
566.8	Flg	577.7	Hu
626.6	F2u	672.7	Gg
701.9	Gu	726.4	F2g
755.8	Gu	770.3	Hu
779.8	Hg	810.9	Flg
926.6	F2g	958.3	Gu
1019.0	F2u	1084.3	Au
1160.0	Hg	1173.5	Gg
1289.7	Hu	1309.4	F2u
1374.1	Flu	1398.5	Hg
1463.8	F2g	1578.5	Hu
1590.0	Gg	1620.4	Gu
1627.4	Ag	1655.3	Flu
1665.1	F2g	1688.2	Hg
1720.1	F2u	1764.4	Gu
1765.4	Gg	1830.0	Hu
1830.7	Hg		

Table 1. Frequencies in cm^{-1} and their degeneracies for the vibrational modes of Buckminsterfullerene. $k_1 = 1.1 \times 10^6$, $k_2 = 1.0 \times 10^6$, $k_3 = 1.0 \times 10^5$ and $k_4 = 1.0 \times 10^5$ dynes/cm.

FIGURE CAPTIONS

- Figure 1. The C_{60} cluster, better known as Buckminsterfullerene, is a truncated icosahedron with 20 hexagonal faces and 12 pentagonal faces.
- Figure 2. A diagram of the local coordinate system used in the calculation.





TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

END

4-87

DTIC